



United Nations Statistics Division

Evaluation of Child Mortality Data from Population Censuses

United Nations Statistics Division



Outline

1. Life tables

- a) Constructing empirical life tables
- b) Model life tables

2. Survival of children ever born (Brass type estimates)

- a) Information required
- b) Checking data quality
- c) Data evaluation using MortPak
- d) Assumptions, violations, and assessing quality of estimates



Life tables



Life tables

- Contain several functions that represent the effects of mortality on a population
 - Life expectancy, age-specific mortality rates, probability of dying by age x
- **Cohort life tables** trace the experience of a single birth cohort (e.g. all those born in 1950)
 - Have to wait for entire cohort to die to have full data
- **Period life tables** use a *synthetic cohort* to represent prevailing mortality conditions at present time
 - As if a cohort lived whole life under current mortality conditions



The period life table – Example (Preston et al. 2001)

Age x	${}_nN_x$	${}_nD_x$	${}_nm_x$	${}_na_x$	${}_nq_x$	${}_nPx$	l_x	${}_nd_x$	${}_nL_x$	T_x	e_x^0
0	47,925	419	0.008743	0.068	0.008672	0.991328	100,000	867	99,192	7,288,901	72.889
1	189,127	70	0.000370	1.626	0.001479	0.998521	99,133	147	396,183	7,189,709	72.526
5	234,793	36	0.000153	2.500	0.000766	0.999234	98,986	76	494,741	6,793,526	68.631
10	238,790	46	0.000193	3.143	0.000963	0.999037	98,910	95	494,375	6,298,785	63.682
15	254,996	249	0.000976	2.724	0.004872	0.995128	98,815	481	492,980	5,804,410	58.740
20	326,831	420	0.001285	2.520	0.006405	0.993595	98,334	630	490,106	5,311,431	54.014
25	355,086	403	0.001135	2.481	0.005659	0.994341	97,704	553	487,127	4,821,324	49.346
30	324,222	441	0.001360	2.601	0.006779	0.993221	97,151	659	484,175	4,334,198	44.613
35	269,963	508	0.001882	2.701	0.009368	0.990632	96,492	904	480,384	3,850,023	39.900
40	261,971	769	0.002935	2.663	0.014577	0.985423	95,588	1,393	474,686	3,369,639	35.252
45	238,011	1,154	0.004849	2.698	0.023975	0.976025	94,195	2,258	465,777	2,894,953	30.734
50	261,612	1,866	0.007133	2.676	0.035082	0.964918	91,937	3,225	452,188	2,429,176	26.422
55	181,385	2,043	0.011263	2.645	0.054861	0.945139	88,711	4,867	432,096	1,976,988	22.286
60	187,962	3,496	0.018600	2.624	0.089062	0.910938	83,845	7,467	401,480	1,544,893	18.426
65	153,832	4,366	0.028382	2.619	0.132925	0.867075	76,377	10,152	357,713	1,143,412	14.971
70	105,169	4,337	0.041238	2.593	0.187573	0.812427	66,225	12,422	301,224	785,699	11.864
75	73,694	5,279	0.071634	2.518	0.304102	0.695898	53,803	16,362	228,404	484,475	9.005
80	57,512	6,460	0.112324	2.423	0.435548	0.564452	37,441	16,307	145,182	256,070	6.839
85	32,248	6,146	0.190585	5.247	1.000000	0.000000	21,134	21,134	110,889	110,889	5.247

Data source: United Nations, 1994.



Calculating the period life table

${}_nM_x$ = age-specific period mortality rate

*# deaths among those aged x to $x+n$
mid-period population aged x to $x+n$*

${}_nq_x$ = probability of dying within next n
years for those who reach age x

l_x = number of people from the original
cohort who live to their x^{th} birthday

*Survivors at age $x+n$ =
survivors at age x * $(1 - {}_nq_x)$*



${}_nL_x$ = number of person-years lived
between exact ages x and $x+n$

e_x = life expectancy at age x = (average
number of years that people have left
to live when they are at age x)

*Sum of all ${}_nL_x$ from age x to
maximum age $\div l_x$*



Data checks: does the life table make sense?



Source: Swedish females, 1895 vs 1995, *Demography*, Preston et. al. 2001



Example, Sudan, 1993 Census, Men - using MortPak LIFTB

MORTPAK FOR WINDOWS - [Selected application is LIFTB (Untitled1)]

File Edit View Application Run Chart Window Help

Input File Name: C:\Program Files (x86)\MORTPAK4\Untitled.MPL
When last updated: 15 October 2014

Construction of a life table.

Age Group	m(x,n)	Age	m(x,n)	q(x,n)	l(x)	d(x,n)	L(x,n)	S(x,n)	T(x)	e(x)	a(x,n)
0 - 1	0.04608	0	0.04608	0.04438	100000.	4438.	96317.	0.94428	5219304.	52.193	0.170
1 - 5	0.00689	1	0.00689	0.02710	95562.	2589.	375823.	0.97592	5122987.	53.609	1.519
5 - 10	0.00355	5	0.00355	0.01759	92972.	1636.	460772.	0.98032	4747164.	51.060	2.500
10 - 15	0.00441	10	0.00441	0.02181	91337.	1992.	451703.	0.97570	4286392.	48.930	2.500
15 - 20	0.00556	15	0.00556	0.02743	89345.	2450.	440726.	0.97134	3834689.	42.920	2.553
20 - 25	0.00601	20	0.00601	0.02961	86894.	2573.	428097.	0.96904	3393963.	39.059	2.523
25 - 30	0.00658	25	0.00658	0.03237	84321.	2730.	414844.	0.96678	2965866.	35.173	2.523
30 - 35	0.00716	30	0.00716	0.03520	81592.	2872.	401064.	0.95549	2551022.	31.266	2.599
35 - 40	0.01139	35	0.01139	0.05545	78720.	4365.	383212.	0.93797	2149958.	27.311	2.620
40 - 45	0.01428	40	0.01428	0.06903	74355.	5133.	359440.	0.91531	1766746.	23.761	2.597
45 - 50	0.02090	45	0.02090	0.09933	69222.	6876.	329001.	0.90691	1407305.	20.330	2.512
50 - 55	0.01860	50	0.01860	0.08902	62346.	5550.	298375.	0.86904	1078304.	17.295	2.593
55 - 60	0.03941	55	0.03941	0.17992	56796.	10219.	259299.	0.81543	779929.	13.732	2.585
60 - 65	0.04139	60	0.04139	0.18789	46578.	8752.	211440.	0.75638	520631.	11.178	2.549
65 - 70	0.07553	65	0.07553	0.31934	37826.	12079.	159929.	0.60444	309190.	8.174	2.583
70 - 75	0.13096	70	0.13096	0.49170	25747.	12860.	96668.	0.42228	149261.	5.797	2.467
75 - 80		75	0.21849	0.68152	13087.	8919.	40821.	0.22383	52593.	4.019	2.240
80 - 85		80	0.35405	...	4168.	4168.	11772.	...	11772.	2.824	2.824
85 - 90											
90 - 95											
95 - 100											

First entry of S(x,n) is for survivorship of 5 cohorts of birth to age group 0-4 = L(0,5) / 500000
Second entry of S(x,n) is for S(0,5) = L(5,5) / L(0,5)
Last entry of S(x,n) is S(75+,5) = T(80) / T(75)

Data source: 1993 Census Report, Table B5 and D3.



Model life tables


- Represent expected age patterns of mortality
- Created to estimate demographic parameters for countries with limited data
- Built on empirical studies of age-specific mortality patterns in the past
- Two groups of model life tables:
 - Coale-Demeny (1983): based on European populations
 - North, South, East and West European models
 - West only model based on some non-European life tables
 - United Nations (1982): based on developing countries
 - Latin American, Chilean, South Asian, Far Eastern, General



Model life tables (2)

1. Age-specific shape of mortality – relative probabilities of dying at different ages

Figure 4. Relationship between infant mortality, $q(1)$, and child mortality, ${}_4q_1$, in the United Nations mortality models





Source: *Step by step guide to the estimation of child mortality*, 1990, United Nations



Model life tables (3)

1. Age-specific shape of mortality – relative probabilities of dying at different ages

Deviation of UN Model Life Tables from Coale-Demeny Model West, Females



Source: United Nations (1982), *Model Life tables for Developing Countries*.



Model life tables (3)

2. Level of mortality – each model has several different levels that correspond with a different life expectancies at birth (e0)

United Nations Model Life Tables — Males

Latin American Pattern

AGE	M(X)	Q(X)	I(X)	D(X)	L(X)	T(X)	E(X)	A(X)
0	.23669	.20429	100000	20429	86313	3500000	35.000	0.330
1	.04672	.16631	79571	13234	283241	3413687	42.901	1.352
5	.00982	.04790	66337	3178	323742	3130446	47.190	2.500
10	.00511	.02522	63160	1593	311817	2806704	44.438	2.500
15	.00697	.03427	61567	2110	302841	2494887	40.523	2.633
20	.01036	.05051	59457	3003	290037	2192046	36.868	2.586
25	.01169	.05679	56454	3206	274346	1902009	33.691	2.528
30	.01352	.06449	53248	3434	257753	1627663	30.567	2.528
35	.01528	.07363	49814	3668	239996	1369910	27.500	2.529
40	.01757	.08418	46146	3885	221132	1129914	24.485	2.531
45	.02092	.09948	42262	4204	200930	908782	21.504	2.538
50	.02517	.11849	38058	4509	179185	707852	18.599	2.542
55	.03225	.14939	33548	5012	155420	528667	15.758	2.543
60	.04241	.19205	28537	5480	129217	373247	13.080	2.520
65	.06056	.26327	23056	6070	100230	244030	10.584	2.461
70	.08574	.35208	16986	5980	69747	143800	8.466	2.386
75	.11840	.45210	11006	4976	42023	74053	6.729	2.295
80	.16226	.56382	6030	3400	20953	32030	5.312	2.295
85	.23745	*****	2630	2630	11077	11077	4.211	4.211

AGE	M(X)	Q(X)	I(X)	D(X)	L(X)	T(X)	E(X)	A(X)
0	.22881	.19840	100000	19840	86707	3599999	36.000	0.330
1	.04434	.15871	80160	12723	286952	3513291	43.828	1.352
5	.00933	.04560	67438	3075	329502	3226339	47.842	2.500
10	.00487	.02408	64363	1550	317940	2896838	45.008	2.500
15	.00667	.03281	62813	2061	309189	2578898	41.057	2.634
20	.00992	.04843	60752	2942	296662	2269709	37.360	2.588
25	.01120	.05451	57810	3151	281263	1973048	34.130	2.529
30	.01277	.06192	54658	3384	264933	1691784	30.952	2.530
35	.01470	.07093	51274	3637	247381	1426852	27.828	2.528
40	.01696	.08140	47637	3878	228615	1179470	24.759	2.532
45	.02029	.09663	43759	4228	208371	950856	21.729	2.535
50	.02452	.11564	39531	4571	186413	742484	18.782	2.541
55	.03156	.14644	34960	5119	162227	556071	15.906	2.545
60	.04164	.18889	29840	5637	135367	393844	13.198	2.546
65	.05958	.25961	24204	6284	105456	258477	10.679	2.523
70	.08453	.34804	17920	6237	73786	153022	8.539	2.464
75	.11698	.44810	11683	5235	44753	79236	6.782	2.390
80	.16076	.56044	6448	3614	22479	34483	5.348	2.299
85	.23611	*****	2834	2834	12004	12004	4.235	4.235

Source: United Nations (1982), *Model Life tables for Developing Countries*.



Survival of children ever born

Indirect estimation of child mortality



Mortality estimates from population censuses: Introduction

- ❑ A group of questions can be used to obtain mortality data in a census
- ❑ Two distinctions:
 - a) Level and trend of mortality vs age pattern of mortality
 - Survival of children ever born: level and trend of mortality
 - Household deaths: age pattern of mortality
 - b) Deaths of younger persons vs. deaths of adults
 - Younger persons: survival of children ever born
 - Adults: household deaths
- ❑ All approaches are to supplement death registration data, not to replace it.



Quick review - children ever born data

- ❑ Have been used for the past 50 years to collect data on **infant and child mortality**
- ❑ For every woman the following information is collected:
 - a) the total number of female children she has had in her lifetime.
 - b) the total number of male children she has had in her lifetime.
 - c) the number of female children who are surviving
 - d) the number of male children who are surviving



Survival of children ever born

- ❑ Ever born – Surviving = Children deceased
- ❑ Children deceased / Ever born = Proportion deceased
- ❑ Life table measures of **infant, child and young adult** mortality may be derived from the proportion of deceased
 - In combination with data on age of mother



Brass type estimates

- ❑ Provide *indirect estimation* of **level and trend** of mortality for about 20 years prior to a census or survey
- ❑ Data required:
 - Number of women by
 - 5 year age group or;
 - Duration of marriage (5 year groups)
 - Total number of children born alive to women in corresponding 5-year groups
 - Total number of children still alive (or deceased) at time of census by corresponding 5-year groups



Brass type estimates (2)

Age group of mother in years	Age group index	Proportion of children dead approximates
15-19	1	q(1)
20-24	2	q(2)
25-29	3	q(3)
30-34	4	q(5)
35-39	5	q(10)
40-44	6	q(15)
45-49	7	q(20)
50-54	8	q(25)
55-59	9	q(30)



Brass type estimation – data checks

Women in the age group should include all women, not only those who respond to CEB/CS questions

>> Important to check in contexts where inappropriate to ask unmarried women about childbearing

BANGLADESH CENSUS 1974 RETROSPECTIVE SURVEY OF FERTILITY AND MORTALITY

AGE GROUP OF WOMEN	TOTAL WOMEN	TOTAL BIRTHS	CHILDREN AT HOME	CHILDREN AWAY	CHILDREN DEAD
TOTAL					
0-14	259 104	6 677	4 866	0	1 811
15-19	2 019 436	1 160 919	921 227	24 327	215 365
20-24	2 521 318	4 901 382	3 820 649	83 349	997 384
25-29	2 573 496	9 085 852	6 927 908	219 989	1 937 955
30-34	2 003 082	9 910 256	7 126 473	522 587	2 261 196
35-39	1 766 100	10 384 001	6 974 267	919 566	2 490 168
40-44	1 473 382	9 164 329	5 472 460	1 276 846	2 415 023
45-49	1 128 791	6 905 673	3 664 328	1 281 801	1 959 544
50-54	1 040 877	5 963 087	2 601 163	1 441 061	1 920 863
55-59	601 625	3 257 428	1 206 148	913 559	1 137 721
60+	1 631 217	8 136 608	2 102 978	2 800 615	3 233 015
N.S.	204	0	0	0	0
TOTAL	17 018 632	68 876 212	40 822 467	9 483 700	18 570 045

Note small number of women in 0-14 age group – unmarried were not included

Source: United Nations (1990), *Step by step guide to the estimation of child mortality*



Brass type estimation – data checks (2)

- ❑ Experience has shown that it is possible to get high quality responses to summary birth histories in any data collection exercise, including censuses
 - > If both CEB and CS are understated, some cancellation of errors will occur
 - > But in practice, reporting of CS is more likely to be complete than reporting of CEB => calculated proportions of deceased children are likely to be too low
- ❑ Make sure trends in children ever born/surviving/deceased are consistent
- ❑ Check for missing data and/or editing



Brass type estimation – data checks (3)

Example: missing or implausible values of CEB and CS data

Table 5.1 Percentage of cases where no editing of children ever born and children surviving data was required, by population group and age group

<i>Age</i>	<i>African</i>		<i>Coloured</i>		<i>Indian/Asian</i>		<i>White</i>	
	<i>CEB</i>	<i>CS</i>	<i>CEB</i>	<i>CS</i>	<i>CEB</i>	<i>CS</i>	<i>CEB</i>	<i>CS</i>
12-14	65.2	34.5	53.5	27.2	61.4	19.7	46.2	22.6
15-19	73.5	44.0	63.7	37.2	68.8	24.6	55.9	28.9
20-24	82.5	62.5	78.5	59.5	79.1	40.9	73.0	44.5
25-29	88.2	75.6	87.6	75.4	88.0	64.0	80.3	62.7
30-34	90.9	81.2	91.2	82.0	92.2	78.0	81.5	71.5
35-39	91.9	83.2	92.6	84.5	93.5	82.0	84.5	78.0
40-44	91.4	83.3	92.5	84.7	93.3	83.0	84.5	82.7
45-49	89.9	82.3	91.3	83.7	91.9	82.6	90.4	82.2

80.3 per cent need to be edited

Source: Dorrington & Moultrie (2001).



Brass type estimation – data checks (4)

Turkey, 2000						
Age group of women	Total women	Total CEB	Average CEB	Total CS	Average children deceased (CD)	Proportion deceased (CEB-CS)/CEB
15 - 19	3518257	294628	0.08	281296	0.003789	0.045
20 - 24	3263432	2078364	0.64	1991445	0.026634	0.042
25 - 29	2918825	4522719	1.55	4312404	0.072055	0.047
30 - 34	2457285	5700038	2.32	5395143	0.124078	0.053
35 - 39	2400808	7036619	2.93	6563946	0.196881	0.067
40 - 44	1985225	6707033	3.38	6131544	0.289886	0.086
45 - 49	1658012	6394157	3.86	5722904	0.404854	0.105

Average CEB should be realistic given country TFR and typical ages at childbearing

Unless fertility or child mortality are increasing, average CD should increase with age group

Unless fertility has been rising, average CEB should increase with age group

Source: Tabulated using data from *Demographic Yearbook*



Brass type estimation – data checks (5)

- ❑ Check sex ratio at birth implied by the CEB data for different mother age groups if gender is disaggregated (from age & sex structure)
 - Is it plausible?
 - Can help to identify underreporting of female births
- ❑ Is proportion of children surviving/deceased plausible?
 - Compare with other sources on child mortality



A rapid assessment of CEB/CS data: Sudan, 1993 census (1)

Age group	CEB	CS	CS/CEB
15 - 19	144,200	124,924	0.866
20 - 24	835,617	727,299	0.870
25 - 29	2,178,788	1,892,963	0.869
30 - 34	2,267,707	1,953,583	0.861
35 - 39	3,335,146	2,841,547	0.852
40 - 44	2,399,753	2,003,693	0.835
45 - 49	2,249,857	1,857,531	0.826



A rapid assessment of CEB/CS data: Sudan, 1993 census (2)

- ❑ Proportion deceased for the 30-34 age group =
 $(1-0.861)=0.139$
 - Proportion of children deceased born to mothers of 30-34 years of age approximates $q(5)$, the proportion of children born who die before their 5th birthday, about 7 years before data collection

- ❑ Compare with other estimates, e.g. UN Population Division estimates of under-5 mortality
 - 1993 census 'quick' estimates of under-5 child mortality = 139 per 1000 for 1986
 - UN Pop Division = 137 per 1000 (for 1985-1990 period)
 - IGME = 134.6 per 1000 (for 1986.5)



UN Population Division: World Population Prospects

<http://esa.un.org/wpp/Excel-Data/mortality.htm>

United Nations, Department of Economic and Social Affairs
Population Division, Population Estimates and Projections Section

United Nations | DESA | Population Division | About Us | Publications | Meetings | Contact

Home
Publications
Frequently Asked Questions
PowerPoint Presentations
Data
Tables in EXCEL-Format
Population
Fertility
Mortality
Migration
Interpolated Data
Data in ASCII-Format
Extended Data Set
On-line Database
Population

World Population Prospects: The 2012 Revision

Excel Tables - Mortality Data

Topic	Data File	Description
Infant and children	Infant Mortality Rate (IMR)	Probability of dying between birth and exact age 1. It is expressed as average annual deaths per 1,000 births.
Infant and children	Under-five Mortality (5q0)	Probability of dying between birth and exact age 5. It is expressed as average annual deaths per 1,000 births.
Overall	Crude Death Rate (CDR)	Number of deaths over a given period divided by the person-years lived by the population over that period. It is expressed as average annual number of deaths per 1,000 population.
Overall	Deaths - Both Sexes	Number of deaths over a given period. Refers to five-year periods running from 1 July to 30 June of the initial and final years. Data are presented in thousands.
Overall	Deaths - Male	Number of male deaths over a given period. Refers to five-year periods running from 1 July to 30 June of the initial and final years. Data are presented in thousands.



IGME: Child mortality estimates

<http://www.childmortality.org>

Child Mortality Estimates
CME Info

Home Map Country Data Compare

Area Selection List

- Sudan
- South Sudan
- Sudan
- Sudan pre secession


Cancel

CME Info is a database containing the latest child mortality estimates based on the research of the UN Inter-agency Group for Child Mortality Estimation.

NEW UPDATES: LATEST RELEASE
16 September 2014
The UN Inter-agency Group for Child Mortality Estimation released the latest estimates on child mortality.



IGME: Child mortality estimates (www.childmortality.org)





Brass type estimation with MortPak **QFIVE**

- ❑ Calculate the sex ratio at birth
 - If not available, can use standard 1.05
- ❑ Calculate the mean age of childbearing (**only for UN model life tables**)

$$M = \frac{17.5 \cdot B_{15-19} + 22.5 \cdot B_{20-24} + \dots + 47.5 \cdot B_{45-49}}{B_{15-19} + B_{20-24} + \dots + B_{45-49}}$$

where $B_{x, x+n}$ = Births in past year to women age x to $x+n$



Brass type estimation with QFIVE in MORTPAK Sudan – 1993 Census

MORTPAK FOR WINDOWS - [Selected application is QFIVE (Untitled1)]

File Edit View Application Run Chart Window Help

Input File Name: C:\Program Files (x86)\MORTPAK4\Untitled.MPL
When last updated: 15 October 2014

Estimates of infant mortality and under 5 mortality by applying the two versions of the Brass method: the True

TITLE: SUDAN, 1993 Census

Month: April
Year: 1993
Sex: Both sexes
Sex Ratio at Birth:
Mean Age at Childbearing:
Data Definition:

Select type of input based on data available

Age Group of Woman	Not defined above	Number of children ever born, number of children surviving and number of women
15 - 20	0.1270	0.1101
20 - 25	0.8963	0.7801
25 - 30	2.3742	2.0627
30 - 35	3.8170	3.2882
35 - 40	5.3358	4.5461
40 - 45	5.8607	4.8935
45 - 50	6.1806	5.1028



Brass output with QFIVE in MORTPAK

9 Models:
5 United Nations; 4 Coale-Demeny

Indicators:

- $q(0)$
- $q(1, 4)$
- $q(5)$
- $e(0)$


Age Group of Woman	United Nations Models (Palloni-Helgman Equations)										Coale-Demeny Models (Trussell Equations)							
	Latin American		Chilean		South Asian		Far East		General		West		North		East		South	
	Ref. Date	q(x)	Ref. Date	q(x)	Ref. Date	q(x)	Ref. Date	q(x)	Ref. Date	q(x)	Ref. Date	q(x)	Ref. Date	q(x)	Ref. Date	q(x)	Ref. Date	q(x)
Infant mortality rate (probability of dying between ages 0 and 1): $q(1)$																		
15 - 20	1992.2	0.130	1992.0	0.145	1992.2	0.130	1992.1	0.132	1992.1	0.131	1992.2	0.139	1992.2	0.135	1992.2	0.141	1992.2	0.131
20 - 25	1991.1	0.110	1991.0	0.127	1991.0	0.111	1991.0	0.112	1991.0	0.112	1991.1	0.115	1991.1	0.105	1991.1	0.121	1991.1	0.111
25 - 30	1989.7	0.097	1989.6	0.116	1989.7	0.099	1989.6	0.100	1989.7	0.100	1989.6	0.103	1989.7	0.092	1989.5	0.111	1989.6	0.102
30 - 35	1988.0	0.092	1987.8	0.114	1988.0	0.095	1987.9	0.096	1988.0	0.095	1987.7	0.100	1987.9	0.089	1987.6	0.110	1987.8	0.100
35 - 40	1986.0	0.091	1985.7	0.115	1985.9	0.095	1985.9	0.093	1986.0	0.094	1985.6	0.100	1985.9	0.087	1985.5	0.111	1985.6	0.102
40 - 45	1983.6	0.093	1983.2	0.120	1983.4	0.100	1983.5	0.094	1983.5	0.096	1983.2	0.103	1983.6	0.088	1983.0	0.117	1983.2	0.105
45 - 50	1980.3	0.093	1979.8	0.119	1979.9	0.101	1980.4	0.090	1980.3	0.095	1980.2	0.100	1980.8	0.084	1979.7	0.115	1980.1	0.104
Probability of dying between ages 1 and 5: $q(1,4)$																		
15 - 20	1992.2	0.099	1992.0	0.048	1992.2	0.089	1992.1	0.085	1992.1	0.087	1992.2	0.080	1992.2	0.108	1992.2	0.056	1992.2	0.091
20 - 25	1991.1	0.074	1991.0	0.038	1991.0	0.069	1991.0	0.064	1991.0	0.066	1991.1	0.060	1991.1	0.077	1991.1	0.044	1991.1	0.065
25 - 30	1989.7	0.060	1989.6	0.032	1989.7	0.057	1989.6	0.054	1989.7	0.055	1989.6	0.051	1989.7	0.063	1989.5	0.039	1989.6	0.053
30 - 35	1988.0	0.055	1987.8	0.031	1988.0	0.053	1987.9	0.049	1988.0	0.050	1987.7	0.049	1987.9	0.060	1987.6	0.038	1987.8	0.051
35 - 40	1986.0	0.054	1985.7	0.031	1985.9	0.053	1985.9	0.047	1986.0	0.049	1985.6	0.049	1985.9	0.058	1985.5	0.039	1985.6	0.053
40 - 45	1983.6	0.056	1983.2	0.034	1983.4	0.057	1983.5	0.048	1983.5	0.051	1983.2	0.051	1983.6	0.060	1983.0	0.042	1983.2	0.057
45 - 50	1980.3	0.056	1979.8	0.034	1979.9	0.058	1980.4	0.045	1980.3	0.050	1980.2	0.049	1980.8	0.056	1979.7	0.041	1980.1	0.058
Probability of dying by age 5: $q(5)$																		
15 - 20	1992.2	0.217	1992.0	0.186	1992.2	0.208	1992.1	0.205	1992.1	0.207	1992.2	0.208	1992.2	0.229	1992.2	0.189	1992.2	0.210
20 - 25	1991.1	0.176	1991.0	0.160	1991.0	0.173	1991.0	0.169	1991.0	0.171	1991.1	0.168	1991.1	0.174	1991.1	0.160	1991.1	0.169
25 - 30	1989.7	0.151	1989.6	0.145	1989.7	0.150	1989.6	0.149	1989.7	0.149	1989.6	0.149	1989.7	0.150	1989.5	0.146	1989.6	0.150
30 - 35	1988.0	0.142	1987.8	0.141	1988.0	0.143	1987.9	0.140	1988.0	0.141	1987.7	0.145	1987.9	0.143	1987.6	0.144	1987.8	0.147
35 - 40	1986.0	0.140	1985.7	0.142	1985.9	0.143	1985.9	0.136	1986.0	0.139	1985.6	0.144	1985.9	0.140	1985.5	0.146	1985.6	0.149
40 - 45	1983.6	0.143	1983.2	0.150	1983.4	0.151	1983.5	0.138	1983.5	0.142	1983.2	0.149	1983.6	0.143	1983.0	0.153	1983.2	0.157
45 - 50	1980.3	0.143	1979.8	0.149	1979.9	0.153	1980.4	0.131	1980.3	0.140	1980.2	0.144	1980.8	0.136	1979.7	0.151	1980.1	0.155
Life expectancy at birth: $e(0)$																		
15 - 20	1992.2	47.021	1992.0	49.425	1992.2	51.520	1992.1	39.917	1992.1	45.600	1992.2	46.750	1992.2	44.325	1992.2	51.174	1992.2	50.291




How to identify the right model life table (1)

Relationship between mortality risk during the first year of life and between ages 1 - 4

Coale-Demeny Models



United Nations Models




Source: United Nations (1990)



How to identify the right model life table (2)


Comparison of $q(1)$ & $q(1,4)$ estimates with Coale-Demeny models






How to identify the right model life table (3)

Direct estimates of $q(0)$ and $q(1,4)$ from Sudan 1990 DHS and 1993 Census, and the relationships to Coale-Demeny and UN model life tables





Estimated under five and under one mortality over time, Sudan 1993 Census (CD North)




Most recent estimates should be disregarded as estimates based on reports from young mothers tend to be exaggerated



Brass: relationship of mother's age and timing of the under-5 mortality estimates

Bangladesh, 1974 Retrospective Survey of Fertility and Mortality

Figure 7. Under-five mortality, $q(5)$, for both sexes in Bangladesh, estimated using model South and the Trussell version of the Brass method



Source: United Nations (1990)



Brass: Assumptions, violations (1)

1. In any time period, mortality of children does not vary by five-year grouping of mothers
 - This assumption is usually violated for the mother age group 15 – 19, and to a lesser extent for the age group 20 – 24, because children of young mothers are known to have higher risk of mortality
 - Why?
 - First births have higher mortality risk than higher-order births and children of younger mothers are more likely to be first births
 - Youngest mothers tend to be socio-economically disadvantaged

Source: Moultrie et al. (2013)



Brass: Assumptions, violations (2)

2. No correlation exists between mortality risks of children and survival of mothers in the population
 - This is a problem when certain mothers are not captured in the data (because of mortality or migration) whose children might also have higher mortality risk
 - Most common case is countries with high HIV prevalence – results in downward bias in estimates
 - Younger mother age groups (20-24, 25-29) less likely to be biased
 - See for adjustment techniques:
<http://demographicestimation.iussp.org/content/effects-hiv-methods-child-mortality-estimation>

Source: Moultrie et al. (2013)



Brass: Assumptions, violations (3)

3. Population age patterns of fertility and child mortality are adequately represented by the model patterns used in developing the method
 4. Any changes in child mortality in the recent past have been gradual and unidirectional
 5. Cross-sectional average numbers of children ever born by age adequately reflect cohort patterns of childbearing
- Note that when fertility has been changing (falling) rapidly, the Brass method will tend to over-estimate child mortality
 - Variants of the technique grouping mothers by duration of marriage or time since first birth have been developed to address some of these issues

Source: Moultrie et al. (2013)



Brass: $q(5)$ more robust to model life table choice than $q(1)$

Figure 12. Infant and under-five mortality for both sexes in Bangladesh, estimated using the four Coale-Demeny mortality models




Figure 13. Infant and under-five mortality for both sexes in Bangladesh, estimated using the five United Nations mortality models





Figure 14. Range of variation of the possible estimates of infant and under-five mortality for both sexes in Bangladesh




Source: United Nations (1990)

Source: Tables 16 and 17.




Quality of estimates: Checking multiple sources






Quality of estimates: Comparing age patterns of proportion of children deceased





Quality of estimates: Comparison with existing external sources




IGME estimates

Source: www.childmortality.org



Quality of estimates: Implied life expectancy at birth using Model Life Tables





References

- Feeney, G. (2009), *Rapid Assessment of Census Data on Children Born and Surviving*, available online at: <http://demographer.com/2009-blog/rapid-assessment-of-ceb-and-cs-data/>
- Moultrie, T. et al. (2013), *Tools for Demographic Estimation*, Paris: IUSSP, available online at: <http://demographicestimation.iussp.org/>
- United Nations (1982), *Model Life Tables for Developing Countries*, New York: United Nations, available online at: http://www.un.org/esa/population/publications/Model_Life_Tables/Model_Life_Tables.htm
- United Nations (1983), *Manual X: Indirect Techniques for Demographic Estimation*, New York: United Nations, available online at: http://www.un.org/esa/population/publications/Manual_X/Manual_X.htm
- United Nations (1990), *Step-by-step Guide to the Estimation of Child Mortality*, New York: United Nations, available online at: http://www.un.org/esa/population/techcoop/DemEst/stepguide_childmort/stepguide_childmort.html
- United Nations Population Division (2012) *Updated UN Model Life Tables*, New York: United Nations, available online at: <http://esa.un.org/unpd/wpp/Model-Life-Tables/download-page.html>
- United Nations Population Division (2013) *World Population Prospects: The 2012 Revision*, New York: United Nations, available online at: <http://esa.un.org/wpp/>